

## **Germinal testicular tumour metastatic exclusively to the spleen**

*Afef Khanfir<sup>1</sup>; Ghassen Marrekchi<sup>1</sup>; Jibene Feki<sup>1</sup>; Salona Makni Krichen<sup>2</sup>; Salah Boujelben<sup>3</sup>; Tabia Boudawara<sup>2</sup>; Jamel Mnif<sup>4</sup>; Mohamed Issam Beyrouti<sup>3</sup>; Mounir Frikha<sup>1</sup>*

*(1) Medical oncology department, Habib Bourguiba hospital, 3029 Sfax-Tunisia*

*(2) Cytology and Histopathology department, Habib Bourguiba hospital, 3029 Sfax-Tunisia*

*(3) Visceral surgery department, Habib Bourguiba hospital, 3029 Sfax-Tunisia*

*(4) Radiology department, Habib Bourguiba hospital, 3029 Sfax-Tunisia*

PAJO, June 2013, 6(2): 34-36

### **Abstract**

Non seminomatous germ cell testicular cancer (NSGTC) often metastasizes to lung and lymph nodes. Spleen metastases are rare. We present an original case of a 27 year-old man with bilateral testicular ectopia history in childhood, presenting intraabdominal mixed non seminomatous germ cell cancer (embryonal carcinoma and Yolk sac tumor) with only spleen nodules in the pre chemotherapy abdominal CT scan. Metastases to spleen were confirmed by splenectomy at the end of chemotherapy; they had the same histology than the primitive tumor.